

Aquatic Mammals Key Word Index

Volumes 41.1-41.44 (Alphabetical)

acoustics

Simard et al., Dolphin Distribution on the West Florida Shelf Using Visual Surveys and Passive Acoustic Monitoring, 41.2, 167-187.

active acoustic space

Miksis-Olds et al., The Impact of Ocean Sound Dynamics on Estimates of Signal Detection Range, 41.4, 444-454.

activity budget

Vermeulen et al., Diurnal and Seasonal Variation in the Behaviour of Bottlenose Dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina, 41.3, 272-283.

Alaska

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.

Aleutian Islands

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

annual consumption

Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.

anthropogenic activity

Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

anthropogenic noise

—Dunlop et al., The Behavioural Response of Humpback Whales (*Megaptera novaeangliae*) to a 20 Cubic Inch Air Gun, 41.4, 412-433.

—von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.

anthropogenic sound

—Baird et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Hawai'i Region, 41.1, 54-64.

—Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.

—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.

—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.

antibiotic

Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.

Arctic

—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

arctic biology

Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.

Atlantic spotted dolphin (*Stenella frontalis*)

Simard et al., Dolphin Distribution on the West Florida Shelf Using Visual Surveys and Passive Acoustic Monitoring, 41.2, 167-187.

bacteria

Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.

baleen whales

Dunlop et al., The Behavioural Response of Humpback Whales (*Megaptera novaeangliae*) to a 20 Cubic Inch Air Gun, 41.4, 412-433.

behavior

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps with and without Side Bands, 41.4, 400-411.

—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.

behavioural response

—Dunlop et al., The Behavioural Response of Humpback Whales (*Megaptera novaeangliae*) to a 20 Cubic Inch Air Gun, 41.4, 412-433.

—Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.

beluga (*Delphinapterus leucas*)

—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.

Bering Sea

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.

—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

BIA

Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

bones

Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.

bottlenose dolphin (*Tursiops truncatus*)

—Blasi et al., Influence of Trammel Nets on the Behaviour and Spatial Distribution of Bottlenose Dolphins (*Tursiops truncatus*) in the Aeolian Archipelago, Southern Italy, 41.3, 295-310.

—Fiorucci et al., Reference Baseline Data for Gastric Cytology in Healthy Bottlenose Dolphins (*Tursiops truncatus*) Under Human Care, 41.3, 345-350.

- Fiorucci et al., Determination of the Main Reference Values in Ultrasound Examination of the Gastrointestinal Tract in Clinically Healthy Bottlenose Dolphins (*Tursiops truncatus*), 41.3, 284-294.
- Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.
- LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.
- Simard et al., Dolphin Distribution on the West Florida Shelf Using Visual Surveys and Passive Acoustic Monitoring, 41.2, 167-187.
- bowhead whale (*Balaena mysticetus*)
- Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
- Bryde's whale (*Balaenoptera edeni*)
- LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.
- California
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- California sea lion (*Zalophus californianus*)
- Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.
- capybaras (*Hydrochoerus hydrochaeris*)
- Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.
- cetaceans
- Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.
- cetacean detection, impacts
- Paiva et al., Reduced Detection of Indo-Pacific Bottlenose Dolphins (*Tursiops aduncus*) in an Inner Harbour Channel During Pile Driving Activities, 41.4, 455-468.
- CetMap
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- chronic shedding
- Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.
- clicks
- Rasmussen et al., Buzzes and High-Frequency Clicks Recorded from Narwhals (*Monodon monoceros*) at Their Wintering Ground, 41.3, 256-264.
- coastal
- Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.
- coastal waters
- Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- conservation
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- controlled exposure experiment
- Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- critical ratio
- Erbe, The Maskogram: A Tool to Illustrate Zones of Masking, 41.4, 434-443.
- distribution
- Simard et al., Dolphin Distribution on the West Florida Shelf Using Visual Surveys and Passive Acoustic Monitoring, 41.2, 167-187.
- distribution tendency
- Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- disturbance
- Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
- diversity
- Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.
- Dutch Continental Shelf
- von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.
- East Coast
- LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
- echolocation
- Rasmussen et al., Buzzes and High-Frequency Clicks Recorded from Narwhals (*Monodon monoceros*) at Their Wintering Ground, 41.3, 256-264.
- endoscopy
- Fiorucci et al., Reference Baseline Data for Gastric Cytology in Healthy Bottlenose Dolphins (*Tursiops truncatus*) Under Human Care, 41.3, 345-350.
- EU MSFD
- Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- Europe
- Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- feeding area
- Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
- Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
- Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
- LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
- feeding behaviour
- Blasi et al., Influence of Trammel Nets on the Behaviour and Spatial Distribution of Bottlenose Dolphins (*Tursiops truncatus*) in the Aeolian Archipelago, Southern Italy, 41.3, 295-310.
- feeding ecology
- Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.

- fin whale (*Balaenoptera physalus*)
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
- fisheries sonar
—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps With and Without Side Bands, 41.4, 400-411.
—Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
- foraging
Rasmussen et al., Buzzes and High-Frequency Clicks Recorded from Narwhals (*Monodon monoceros*) at Their Wintering Ground, 41.3, 256-264.
- foraging strategy
Vermeulen et al., Diurnal and Seasonal Variation in the Behaviour of Bottlenose Dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina, 41.3, 272-283.
- gadoids
Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.
- gastric chambers
Fiorucci et al., Determination of the Main Reference Values in Ultrasound Examination of the Gastrointestinal Tract in Clinically Healthy Bottlenose Dolphins (*Tursiops truncatus*), 41.3, 284-294.
- Giardia
Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.
- gray whale (*Eschrichtius robustus*)
—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
- group size
—Blasi et al., Influence of Trammel Nets on the Behaviour and Spatial Distribution of Bottlenose Dolphins (*Tursiops truncatus*) in the Aeolian Archipelago, Southern Italy, 41.3, 295-310.
—Vermeulen et al., Diurnal and Seasonal Variation in the Behaviour of Bottlenose Dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina, 41.3, 272-283.
- Gulf of Alaska
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- Gulf of Mexico
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.
—Simard et al., Dolphin Distribution on the West Florida Shelf Using Visual Surveys and Passive Acoustic Monitoring, 41.2, 167-187.
- habitat
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- habitat characteristic
Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- habitat use
Blasi et al., Influence of Trammel Nets on the Behaviour and Spatial Distribution of Bottlenose Dolphins (*Tursiops truncatus*) in the Aeolian Archipelago, Southern Italy, 41.3, 295-310.
- harbor porpoise (*Phocoena phocoena*)
—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
—von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.
- harbor seal (*Phoca vitulina*)
Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
- Hawai'i
Baird et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Hawai'i Region, 41.1, 54-64.
- Hawaiian Islands
Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- hearing loss
von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.
- humpback whale (*Megaptera novaeangliae*)
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
—Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- Indo-Pacific Bottlenose Dolphin (*Tursiops aduncus*)
—Paiva et al., Reduced Detection of Indo-Pacific Bottlenose Dolphins (*Tursiops aduncus*) in an Inner Harbour Channel During Pile Driving Activities, 41.4, 455-468.
- Indo-Pacific humpback dolphin (*Sousa chinensis*)
—Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- intraspecific competition
Vermeulen et al., Diurnal and Seasonal Variation in the Behaviour of Bottlenose Dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina, 41.3, 272-283.
- legislation
Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- Leptospira interrogans*
Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.
- leptospirosis
Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.
- mammal
Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.

- management
Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- marine acoustics
von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.
- marine ecology
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- marine mammals
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- masking
Erbe, The Maskogram: A Tool to Illustrate Zones of Masking, 41.4, 434-443.
- migratory corridor
—Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
- minke whale (*Balaenoptera acutorostrata*)
Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- mitigation
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps with and without Side Bands, 41.4, 400-411.
- mitigation efficiency
Leaper et al., A Simulation Framework to Evaluate the Efficiency of Using Visual Observers to Reduce the Risk of Injury from Loud Sound Sources, 41.4, 375-387.
- MMO
Leaper et al., A Simulation Framework to Evaluate the Efficiency of Using Visual Observers to Reduce the Risk of Injury from Loud Sound Sources, 41.4, 375-387.
- MMPA
Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- narwhals (*Monodon monoceros*)
Rasmussen et al., Buzzes and High-Frequency Clicks Recorded from Narwhals (*Monodon monoceros*) at Their Wintering Ground, 41.3, 256-264.
- naval sonar
—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps with and without Side Bands, 41.4, 400-411.
—Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
—Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- noise
—Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
—Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
- noise impact
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps with and without Side Bands, 41.4, 400-411.
- noise level
Miksis-Olds et al., The Impact of Ocean Sound Dynamics on Estimates of Signal Detection Range, 41.4, 444-454.
- noise pollution
Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- North Pacific Ocean
—Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- North Pacific right whale (*Eubalaena japonica*)
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
- northern bottlenose whale (*Hyperoodon ampullatus*)
Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- Northwest Atlantic Ocean
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
- odontocete
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz Sonar Down-Sweeps with and without Side Bands, 41.4, 400-411.
- offshore
Hernandez-Milian et al., Insights into the Trophic Ecology of Bottlenose Dolphins (*Tursiops truncatus*) in Irish Waters, 41.2, 226-239.
- Oregon
Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.
- parasites
Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.
- Phocid
Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.
- photo-ID
Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- pile driving
Paiva et al., Reduced Detection of Indo-Pacific Bottlenose Dolphins (*Tursiops aduncus*) in an Inner Harbour Channel During Pile Driving Activities, 41.4, 455-468.
- pinnipeds
Kastelein et al., Behavioral Responses of Harbor Seals (*Phoca vitulina*) to Sonar Signals in the 25-kHz Range, 41.4, 388-399.

- policy
Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- population size
Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- port construction
Paiva et al., Reduced Detection of Indo-Pacific Bottlenose Dolphins (*Tursiops aduncus*) in an Inner Harbour Channel During Pile Driving Activities, 41.4, 455-468.
- power spectrum model of masking
Erbe, The Maskogram: A Tool to Illustrate Zones of Masking, 41.4, 434-443.
- reference intervals
Fiorucci et al., Reference Baseline Data for Gastric Cytology in Healthy Bottlenose Dolphins (*Tursiops truncatus*) Under Human Care, 41.3, 345-350.
- reference values
Fiorucci et al., Determination of the Main Reference Values in Ultrasound Examination of the Gastrointestinal Tract in Clinically Healthy Bottlenose Dolphins (*Tursiops truncatus*), 41.3, 284-294.
- renal disease
Prager et al., Antibiotic Efficacy in Eliminating Leptospirosis in California Sea Lions (*Zalophus californianus*) Stranding with Leptospirosis, 41.2, 203-212.
- reproductive area
—Baird et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Hawai'i Region, 41.1, 54-64.
—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
- resident population
—Baird et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Hawai'i Region, 41.1, 54-64.
—Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – East Coast Region, 41.1, 17-29.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.
- seasonal patterns
Vermeulen et al., Diurnal and Seasonal Variation in the Behaviour of Bottlenose Dolphins (*Tursiops truncatus*) in Bahía San Antonio, Patagonia, Argentina, 41.3, 272-283.
- seismic
Dunlop et al., The Behavioural Response of Humpback Whales (*Megaptera novaeangliae*) to a 20 Cubic Inch Air Gun, 41.4, 412-433.
- seismic survey
Leaper et al., A Simulation Framework to Evaluate the Efficiency of Using Visual Observers to Reduce the Risk of Injury from Loud Sound Sources, 41.4, 375-387.
- sentinels
Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.
- severity scoring
Sivle et al., Severity of Expert-Identified Behavioural Responses of Humpback Whale, Minke Whale, and Northern Bottlenose Whale to Naval Sonar, 41.4, 469-502.
- signal detection
Miksis-Olds et al., The Impact of Ocean Sound Dynamics on Estimates of Signal Detection Range, 41.4, 444-454.
- small and resident population
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Aleutian Islands and Bering Sea Region, 41.1, 79-93.
—Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Alaska Region, 41.1, 65-78.
- soundscape
Miksis-Olds et al., The Impact of Ocean Sound Dynamics on Estimates of Signal Detection Range, 41.4, 444-454.
- species distribution
—Baird et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Hawai'i Region, 41.1, 54-64
—Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
—Clarke et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Arctic Region, 41.1, 94-103.
—LaBrecque et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Gulf of Mexico Region, 41.1, 30-38.
- strandings
Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.
- Thailand
Jutapruet et al., Population Size and Habitat Characteristics of the Indo-Pacific Humpback Dolphin (*Sousa chinensis*) Off Donsak, Surat Thani, Thailand, 41.2, 129-142.
- Toxoplasma gondii*
Obusan et al., Occurrence of Human Pathogenic Bacteria and *Toxoplasma gondii* in Cetaceans Stranded in the Philippines: Providing Clues on Ocean Health Status, 41.2, 149-166.
- trammel nets
Blasi et al., Influence of Trammel Nets on the Behaviour and Spatial Distribution of Bottlenose Dolphins (*Tursiops truncatus*) in the Aeolian Archipelago, Southern Italy, 41.3, 295-310.
- tropical
Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.
- Tursiops truncatus* cytology
Fiorucci et al., Reference Baseline Data for Gastric Cytology in Healthy Bottlenose Dolphins (*Tursiops truncatus*) Under Human Care, 41.3, 345-350.
- U.S.
Dolman & Jasny, Evolution of Marine Noise Pollution Management, 41.4, 357-374.
- U.S. West Coast
Calambokidis et al., Biologically Important Areas for Selected Cetaceans Within U.S. Waters – West Coast Region, 41.1, 39-53.
- ultrasonic sonar
Kastelein et al., Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to 25-kHz FM Sonar Signals, 41.3, 311-326.
- ultrasound
Fiorucci et al., Determination of the Main Reference Values in Ultrasound Examination of the Gastrointestinal Tract in Clinically Healthy Bottlenose Dolphins (*Tursiops truncatus*), 41.3, 284-294.
- underwater explosions
von Benda-Beckmann et al., Assessing the Impact of Underwater Clearance of Unexploded Ordnance on Harbour Porpoises (*Phocoena phocoena*) in the Southern North Sea, 41.4, 503-523.
- underwater noise
—Leaper et al., A Simulation Framework to Evaluate the Efficiency of Using Visual Observers to Reduce the Risk of Injury from Loud Sound Sources, 41.4, 375-387.

—Paiva et al., Reduced Detection of Indo-Pacific Bottlenose Dolphins (*Tursiops aduncus*) in an Inner Harbour Channel During Pile Driving Activities, 41.4, 455-468.

Washington

Ferguson et al., Biologically Important Areas for Cetaceans Within U.S. Waters – Overview and Rationale, 41.1, 2-16.

wildlife

Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.

zone of masking

Erbe, The Maskogram: A Tool to Illustrate Zones of Masking, 41.4, 434-443.

zoonosis

Souza et al., Endoparasite Fauna of Wild Capybaras (*Hydrochoerus hydrochaeris*) (Linnaeus, 1766) from the Upper Paraná River Floodplain, Brazil, 41.2, 213-221.