

Coordinated feeding by Clymene dolphins (*Stenella clymene*) in the Gulf of Mexico

D. Fertl¹, A. J. Schiro² and D. Peake³

¹Minerals Management Service, US Department of the Interior, 1201 Elmwood Park Blvd, New Orleans, LA 70123, USA

²Marine Mammal Research Program, Texas A & M University, 4700 Avenue U, Bldg 303, Galveston, TX 77551, USA

³30 LeBrun Court, Galveston, TX 77551, USA

Little is known of the natural history of the Clymene dolphin (*Stenella clymene*). Available information on its feeding habits is limited to stomach contents of two individuals. One animal that stranded in New Jersey contained squid beaks and small otoliths, mostly myctophids (Perrin *et al.*, 1981); the second specimen had squid beaks and eye-lenses (N. B. Barros and D. K. Odell, Southeast US Stranding Network, unpubl. data). Perrin *et al.* (1981) presumed that Clymene dolphins may take prey largely at night and in deep waters, since myctophids are mesopelagic and most species vertically migrate to the water's surface at night. We report an account of Clymene dolphins feeding in a coordinated manner on schooling fish in the Gulf of Mexico. Our observations are of interest because the feeding episode described here occurred during the daytime, and observations on feeding behaviors by this species have not previously been published.

On 24 September 1996, between 1420 and 1500 h, a group of approximately 30 dolphins was observed feeding in a coordinated manner in the Gulf of Mexico, south of Port O'Connor, Texas (27°12.44 N; 95°17.61 W) in water 1243 m deep. Observations were made from the 19.8 m vessel M/V *Chip XIII*, using handheld 10 × 40 power binoculars. Overcast skies, with a Beaufort sea state of 3, made detailed observations difficult.

The dolphins were observed feeding in an area of 15 m diameter (based on comparison with adult dolphin body lengths of approximately 2 m). The dolphins displayed a negative response (ceasing their coordinated behaviors and moving away) twice when the boat approached within 100 m of the activity; therefore, the boat was kept at an appropriate distance so as to not disrupt the dolphins.

Clymene dolphins are easily confused with spinner dolphins (Jefferson *et al.*, 1993). These dolphins were identified as Clymene dolphins based on the

following criteria: a body size and shape that is more robust than that of the spinner dolphin; a shorter rostrum than that of the spinner; and a dark gray cape that dipped above the eye and below the dorsal fin. From the observation distance, it was not possible to see a distinctive black 'moustache' on the rostrum. No birds were associated with the feeding activity. The fish were approximately 10–15 cm in length and slender; it was not possible to identify the fish species. While the majority of the dolphins worked to keep the fish together in a ball near the surface, individuals appeared to take turns pursuing fish. The containment was very fluid, with many of the surfacings being made counter-clockwise to the fish school. Some dolphins passed through the edge of the aggregation, rolling on their sides in an arc towards the fish. Some dolphins were observed gliding through the fringes, and even the center, of the fish school. They became more active outside the fish school, with some splashing as the animals arced back towards the school. After 40 minutes of observation, the boat resumed its course, while the dolphins continued their activity.

Clymene dolphins have been sighted primarily in deep waters (250–5000 m or deeper) (Mullin *et al.*, 1994; Perrin & Mead, 1994). This report is consistent with the known range of this species in the Gulf of Mexico, where it occurs in water depths ranging from at least 704–3064 m (Mullin *et al.*, 1994).

Using a variety of feeding behaviors, many dolphin species are opportunistic and take advantage of available prey. Opportunistic observations of coordinated feeding on fish schools have been made for a number of delphinids, including killer whales (*Orcinus orca*) (Similä & Ugarte, 1993), dusky dolphins (*Lagenorhynchus obscurus*) (Würsig & Würsig, 1980), Atlantic spotted dolphins (*Stenella frontalis*) (Martin, 1986; Fertl & Würsig, 1995), long-beaked common dolphins (*Delphinus capensis*) (Gallo-Reynoso, 1991), rough-toothed dolphins (*Steno bredanensis*) (Smeenk & Richards,

1995; Steiner, 1995), and bottlenose dolphins (*Tursiops truncatus*) (Bel'kovich *et al.*, 1991; Shane, 1990). It is not known whether this reported observation is typical or atypical of Clymene dolphin feeding habits or behavior. It has been suggested that the Clymene dolphin's diet is similar to that of the spinner dolphin (Perrin *et al.*, 1981). Like the spinner dolphin, the Clymene dolphin probably feeds primarily on mesopelagic fish and squid. Both dolphin species presumably feed opportunistically on epipelagic fish occurring near the surface. For example, spinner dolphins have been observed feeding on flying fish (Würsig *et al.*, 1994).

Fifteen years after its redescription, the Clymene dolphin remains a poorly known cetacean (e.g. Perrin & Mead, 1994; Jefferson *et al.*, 1995; Jefferson, 1996). As noted by Mullin *et al.* (1994), continued monitoring of cetacean distribution and numbers in the Gulf of Mexico will add much needed information to the knowledge of this species.

Acknowledgements

We thank the crew of the *Chip XIII*. N. Barros and D. K. Odell provided unpublished data on stomach contents. T. Jefferson, K. Mullin, D. Weller, B. Würsig, and an anonymous reviewer provided comments on this paper. A Schiro's participation represents Contribution No. 54 of the Marine Mammal Research Program, Texas A & M University at Galveston.

References

- Bel'kovich, V. M., Ivanova, E. E., Yefremenkova, O. V., Kozarovitsky, L. B. & Kharitonov, S. B. (1991) Searching and hunting behavior in the bottlenose dolphin (*Tursiops truncatus*) in the Black Sea. In: K. Pryor & K. Norris (eds) *Dolphin societies: discoveries and puzzles*. pp. 38–67. University of California Press: Berkeley, CA.
- Fertl, D. & Würsig, B. (1995) Coordinated feeding by Atlantic spotted dolphins (*Stenella frontalis*) in the Gulf of Mexico. *Aquatic Mammals* **21**, 3–5.
- Gallo-Reynoso, J. P. (1991) Group behavior of common dolphins (*Delphinus delphis*) during prey capture. *Annales Inst. Biol. Univ. Nac. Auton. Mexico, Ser. Zool.* **62**, 253–262.
- Jefferson, T. A. (1996) Morphology of the Clymene dolphin (*Stenella clymene*) in the northern Gulf of Mexico. *Aquatic Mammals* **22**, 35–43.
- Jefferson, T. A., Leatherwood, S. & Webber, M. A. (1993) Marine mammals of the world: FAO species identification guide. FAO, Rome.
- Jefferson, T. A., Odell, D. K. & Prunier, K. T. (1995) Notes on the biology of the Clymene dolphin (*Stenella clymene*) in the northern Gulf of Mexico. *Marine Mammal Science* **11**, 564–573.
- Martin, A. R. (1986) Feeding association between dolphins and shearwaters around the Azores Islands. *Canadian Journal of Zoology* **64**, 1372–1374.
- Mullin, K. D., Higgins, L. V., Jefferson, T. A. & Hansen, L. J. (1994) Sightings of the Clymene dolphin (*Stenella clymene*) in the Gulf of Mexico. *Marine Mammal Science* **10**, 464–470.
- Perrin, W. F. & Mead, J. G. (1994) Clymene dolphin *Stenella clymene* (Gray, 1846). In: S. H. Ridgway & R. Harrison (eds) *Handbook of Marine Mammals, Volume 5: The first book of dolphins*. pp. 161–171. Academic Press: San Diego, CA.
- Perrin, W. F., Mitchell, E. D., Mead, J. G., Caldwell, D. K. & van Bree, P. J. H. (1981) *Stenella clymene*, a rediscovered tropical dolphin of the Atlantic. *Journal of Mammalogy* **62**, 583–598.
- Shane, S. H. (1990) Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. In: S. Leatherwood & R. R. Reeves (eds) *The bottlenose dolphin*. pp. 541–558. Academic Press: San Diego, CA.
- Similä, T. & Ugarte, F. (1993) Surface and underwater observations of cooperatively feeding killer whales in northern Norway. *Canadian Journal of Zoology* **71**, 1494–1499.
- Smeenk, C. & Richards, H. (1995) Some observations on the behaviour of wild rough-toothed dolphins (*Steno bredanensis*). Abstract, European Association for Aquatic Mammals, 23rd Annual Symposium. 31 March–3 April, 1995, Nuremberg, Germany.
- Steiner, L. (1995) Rough-toothed dolphin, *Steno bredanensis*: a new species record for the Azores, with some notes on behaviour. Arquipélago. *Life and Marine Sciences* **13A**, 125–127.
- Würsig, B. & Würsig, M. (1980) Behavior and ecology of the dusky dolphin, *Lagenorhynchus obscurus*, in the South Atlantic. *US Fishery Bulletin* **77**, 871–890.
- Würsig, B., Wells, R. S. & Norris, K. S. (1994) Food and feeding. In: K. S. Norris, B. Würsig, R. S. Wells & M. Würsig (eds) *The Hawaiian spinner dolphin*. pp. 216–231. University of California Press: Berkeley, CA.