

**Necrocoitus in Common Bottlenose Dolphins
(*Tursiops truncatus*) near Sarasota, Florida**

Supplementary Material

Amber Lea D. Kincaid,¹ Gretchen N. Lovewell,¹
Jason B. Allen,² Kim Bassos-Hull,^{1,2} Jessica L. Blackburn,¹
Rebecca A. Hazelkorn,^{1,3} and Randall S. Wells²

¹Mote Marine Laboratory, 1600 Ken Thompson Parkway, Sarasota, FL 34236, USA
E-mail: aldkincaid@gmail.com

²Chicago Zoological Society's Sarasota Dolphin Research Program,
c/o Mote Marine Laboratory, 1600 Ken Thompson Parkway, Sarasota, FL 34236, USA

³NOAA, National Marine Fisheries Service, 263 13th Avenue S, St. Petersburg, FL 33701, USA

Supplemental Figures




Figure S1. Map of SDRP sighting locations of ULYS, F178, and F188 prior to ULYS's death




Figure S2. Map of SDRP sighting locations of ATNA, BELY, and TNBS prior to ATNA's death




Figure S3. Map of SDRP sighting locations of FHIS, F142, and F276 prior to FHIS's death




Figure S4. Map of SDRP sighting locations of F167, C834, and F182 prior to F167's death